


Journal Pre-proof

Evaluating a Community Health Worker Training Program for Medication Adherence Support

Richard Segal R.Ph., Ph.D. ,
Michelle L. Zeigler Pharm.D. Ph.D., BCACP ,
Jordan L. Wallace PharmD, BCOP ,
Teresa E. Roane Pharm.D., MBA, BCACP ,
Lorraine Mobley R.Ph.,C.Ph.,BCGP , Folakemi T. Odedina Ph.D.



PII: S2773-0654(25)00026-4
DOI: <https://doi.org/10.1016/j.focus.2025.100338>
Reference: FOCUS 100338

To appear in: *AJPM Focus*

Received date: 11 November 2024
Revised date: 15 March 2025
Accepted date: 19 March 2025

Please cite this article as: Richard Segal R.Ph., Ph.D. , Michelle L. Zeigler Pharm.D. Ph.D., BCACP , Jordan L. Wallace PharmD, BCOP , Teresa E. Roane Pharm.D., MBA, BCACP , Lorraine Mobley R.Ph.,C.Ph.,BCGP , Folakemi T. Odedina Ph.D. , Evaluating a Community Health Worker Training Program for Medication Adherence Support, *AJPM Focus* (2025), doi: <https://doi.org/10.1016/j.focus.2025.100338>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2025 Published by Elsevier Inc.
This is an open access article under the CC BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Highlights

- CHWs are well-positioned to address adherence barriers but often lack specific training in improving adherence.
- The training aimed to equip CHWs with skills to address providing medication adherence support.
- The training resulted in high confidence levels in performing key skills related to improving adherence.
- The training has the potential to improve chronic disease management among underserved populations.

Journal Pre-proof

Evaluating a Community Health Worker Training Program for Medication Adherence Support

Authors

Richard Segal, R.Ph., Ph.D.¹ Michelle L. Zeigler, Pharm.D.² Ph.D., BCACP Jordan L. Wallace, PharmD, BCOP³ Teresa E. Roane, Pharm.D., MBA, BCACP⁴ Lorraine Mobley, R.Ph.,C.Ph.,BCGP⁵ Folakemi T. Odedina, Ph.D.⁶

¹Richard Segal, R.Ph., Ph.D.

Professor, Department of Pharmaceutical Outcomes and Policy, College of Pharmacy, University of Florida, 1889 Museum Road, Suite 6300, Gainesville Florida, USA, 32611, segal@cop.ufl.edu

²Michelle L. Zeigler, Pharm.D., Ph.D., BCACP

Medication Management Pharmacist, Aledade, Inc.. Gainesville, FL. At the time of research, she was Clinical Assistant Professor, College of Pharmacy, University of Florida, 1225 Center Drive, Room 3334, Gainesville, Florida, USA, 32610,

³Jordan L. Wallace, PharmD, BCOP

Pediatric Oncology Pharmacy Clinical Specialist, Department of Pharmacy, Golisano Children's Hospital, 9981 S. Healthpark Drive, Fort Myers, Florida, 33908, USA, At the time of research, she was a pharmacy student, College of Pharmacy, University of Florida, Gainesville, Florida, USA, 32610,

⁴Teresa E. Roane, Pharm.D., MBA, BCACP

Clinical Associate Professor and Director of Continuing Education, Department of Pharmacy Education and Practice, College of Pharmacy University of Florida, 1225 Center Drive, Room 3334, Gainesville Florida, USA, 32610,

⁵Lorraine Mobley, R.Ph.,C.Ph.,BCGP

Clinical Lecturer, Department of Pharmacy Education and Practice, College of Pharmacy, University of Florida, 1225 Center Drive, Room 3334, Gainesville Florida, USA, 32610, .

⁶Folakemi T. Odedina, Ph.D.

Cancer Prevention, Survivorship and Care Delivery (CPSCD) Research Program and iCCaRE for Black Men Consortium; Mayo Clinic Comprehensive Cancer Center, Jacksonville, Fl; and Prostate Cancer Transatlantic Consortium (CaPTC), 4500 San Pablo Road S, Jacksonville, Florida, 32224, USA,

Corresponding Author

Richard Segal, R.Ph., Ph.D.

University of Florida

1889 Museum Road

Box 100496

Gainesville FL 32611

segal@ufl.edu

Word Count: 3000

CRedit author statement: **Richard Segal**: Conceptualization, Methodology, Formal Analysis, Investigation, Writing – Original Draft, Writing – Review & Editing, Supervision, Project administration, Funding Acquisition. **Michelle L. Zeigler**: Investigation, Writing – Original Draft, Writing – Review & Editing, **Jordan L. Wallace**: Investigation, Writing – Original Draft, Writing – Review & Editing, Project Administration, **Teresa E. Roane**: Investigation, Writing – Review & Editing, **Lorraine Mobley**: Investigation, Writing – Review & Editing, Folakemi T. Odedina: Conceptualization, Investigation, Writing – Review & Editing.

Declaration of Interest

Conflict of interest statement: None. Richard Segal, Michelle Zeigler, Jordan Wallace, Teresa Roane, Lorraine Mobley, Folake Odeina have no conflict disclosures other than to disclose a funding source for this project from the *State Public Health Collaborative to Improve Cardiovascular Health Outcomes* project and Cooperative Agreement number NU58DP004800, funded by the Centers for Disease Control and Prevention.

The paper's contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services.

This publication was supported (in part) by the *State Public Health Collaborative to Improve Cardiovascular Health Outcomes* project and Cooperative Agreement number NU58DP004800, funded by the Centers for Disease Control and Prevention.

The University of Florida IRB approved a protocol associated with the work described in the manuscript.

The study sponsor did not have any role in study design; collection, analysis, and interpretation of data; writing the report; and the decision to submit the report for publication.

Financial Disclosure: No financial disclosures were reported by the authors of this paper. None of the authors have financial relationships with entities that did not support the study, but that might reasonably be considered to be stakeholders in the overall research topic.

Abstract

Introduction: Medication nonadherence significantly contributes to poor health outcomes and increased healthcare costs, particularly in chronic diseases like hypertension. Community Health Workers (CHWs)

are well-positioned to address adherence barriers but often lack specific training in medication management and particularly in improving medication adherence.. The objective of this study is to evaluate a training program for CHWs focused on medication adherence support for patients with hypertension.

Study Design: We designed a comprehensive training program consisting of pre-recorded lectures and a two-day live session. The live training content included didactic lectures integrated with hands-on practice cases and small group breakout sessions. The program was evaluated using pre- and post-tests assessing knowledge and self-efficacy. A quality improvement process was implemented post-training to support CHWs in their new roles.

Results: 109 CHWs participated in the training. Knowledge scores improved from 28% pre-training to 81% post-training ($p < 0.001$). Self-efficacy scores showed significant improvement across all measured domains.

Conclusion: This structured training program effectively improved CHWs' knowledge and confidence in supporting medication adherence. This approach shows promise for enhancing CHWs' roles in collaborating with healthcare professionals for chronic disease management.

Key Words: Medication adherence, Community Health Workers, Pharmacists, Training program, Marginalized populations, Hypertension

Introduction

Poor adherence to prescribed medications significantly contributes to morbidity, mortality, and increased healthcare costs.^{1,2} Among chronically ill patients, about 50% do not take medications as

prescribed.³⁻⁵ In the United States, medication nonadherence is associated with 4 to 28% of medication-related hospital admissions.⁶⁻⁹ Cardiovascular disease, particularly hypertension, presents a significant challenge regarding medication adherence.¹⁰ Despite the known benefits of antihypertensive, lipid-lowering, and antidiabetic medications in reducing ischemic events, adherence to these therapies is poor,¹¹ with studies reporting that 50 to 80% of patients treated for hypertension are non-adherent to their prescribed regimens.¹²⁻¹⁴ The World Health Organization (WHO) considers this lack of adherence the most significant cause of failure to achieve blood pressure control.⁵

Multiple factors contribute to medication nonadherence, including forgetfulness, complex regimens, insufficient education about medication benefits, concerns about side effects, and medication costs. Health literacy also plays a crucial role, with an estimated 90 million adults in the U.S. having inadequate health literacy, a factor particularly prevalent in minority groups.¹⁵⁻¹⁸ Additionally, patients' health beliefs, previous experiences with therapies, lack of motivation, language barriers, lower socioeconomic status, lack of family or social support, and fewer available resources also affect medication adherence.¹⁹⁻²¹

Pharmacists are well-positioned to evaluate and educate patients about medication adherence. However, a survey of 275 patients indicated that those with limited health literacy were less likely to ask their local pharmacist questions about medications.¹⁸ Pharmacists often face difficulties in improving medication use and addressing adherence barriers, especially in culturally diverse populations where they usually lack shared life experiences. Patients may be more willing to express concerns about their health and medication use practices with individuals with shared life experiences.²²

Community Health Workers (CHWs) have emerged as a promising solution to address these challenges, especially in culturally diverse communities.²³⁻²⁵ CHWs often share similar backgrounds and experiences with the individuals they serve, allowing for a better understanding of the language, culture, and traditions affecting medication use. This shared experience can facilitate uncovering health information that might otherwise go unnoticed by primary care physicians or pharmacists.²² Further, Brownstein et al.²⁶ conducted a systematic review of CHW interventions for hypertension care, highlighting the potential of CHWs in improving blood pressure control.

However, CHWs often lack specific training in medication management and adherence support. To address this gap, we evaluated a targeted training program designed to enhance CHWs' knowledge and skills in collaborating with healthcare professionals in improving medication adherence among patients with hypertension. We previously showed that CHWs who completed this training program were able to improve their clients' medication adherence and blood pressure control,²⁷

Methods

CHW Training Program Description

The training program was developed through a collaboration between the State of Florida Department of Health and Centers for Disease Control and Prevention (CDC) and was approved by the University of Florida IRB. The training focused on medication adherence assessments to supplement other training that CHWs had already completed. It consisted of a two-day live program supplemented with six hours of pre-recorded lectures watched before the in-person program. The live training content included didactic lectures integrated with hands-on practice cases and small group breakout sessions led by two or more clinical pharmacist faculty trainers, including five or six CHWs per group. Small group sessions allow CHWs to maximize their participation in an interactive format, share their previous experiences,

and apply new skills to facilitate interventions related to medication adherence. Two patient case studies were used to teach various medication management skills, with student pharmacists playing the role of simulated patients. The patient case studies are based on real-life scenarios from clinical practice, featuring patients with poorly controlled high blood pressure and facing challenges and barriers to medication adherence due to poor medication practices. The curriculum for the two-day live program format is illustrated in Figure 1.

Study Design

This project aimed to assess the effectiveness of the two-day live format of the CHW training program by evaluating changes in knowledge and self-efficacy changes through pre- and post-tests.

Participants and Study Settings

CHWs were employed by five organizations in Florida, including a large managed care health plan providing care in clinics across large urban counties (45 CHWs), federally qualified community health center in a large urban county (19 CHWs), free healthcare clinic in a large urban city (6 CHWs), county aging organization (3 CHWs), and an organization serving healthcare needs of American Indians and other underrepresented populations in both rural and urban counties (36 CHWs). Supervisors from each organization selected CHWs to participate in the training. CHWs were tasked with managing the care of patients diagnosed with uncontrolled high blood pressure who were prescribed at least one antihypertensive medication associated with prescription refill data showing poor adherence based on a Proportion of Days Covered (PDC) below 0.80.²⁸ The structure of the two-day live training program was uniform for CHW program participants from each of the five organization and each session was delivered exclusively to CHWs from a single organization.

Measures, Data Collection and Statistical Analyses

The program evaluation focused on two variables: knowledge and self-efficacy (see Table 1 for operational definitions). The knowledge assessment included open-ended questions about Medication Therapy Management (MTM) principles and multiple-choice questions on health disparities and cultural competence. The MTM assessment required short answers to 13 open-ended questions, which were graded by student pharmacists using a rubric. This rubric covered concepts related to identifying medication use practices, adherence barriers, and strategies to resolve these barriers. To achieve a high score, CHWs needed to demonstrate a thorough understanding of these concepts. The self-efficacy assessment used a scale from 0 (cannot do at all) to 10 (highly certain can do) across 14 activities/tasks related to MTM services.

A panel of eight pharmacy faculty members and practitioners reviewed the study instruments for face and content validity, assessing clarity, relevance, and comprehensiveness. Feedback was gathered through structured questionnaires, and revisions were made based on the panel's feedback.

Statistical analysis was conducted to evaluate whether the training program improved CHWs' knowledge of MTM principles and self-efficacy. Paired t-tests were performed to determine the significance of improvements within the CHW group from pre- to post-training. To account for multiple comparisons, Bonferroni correction was applied, adjusting the significance level to 0.0038 for MTM assessment questions and 0.002 for self-efficacy questions. For the self-efficacy assessment, we calculated the mean change score for each of the 14 items from pre- to post-test. Additionally, we determined the percentage of trainees reporting post-training scores in each self-efficacy category (low, moderate, or high).

To evaluate the consistency of improvements across the five organizations employing CHWs, we used a linear mixed-effects model. In this model, pre- and post-program scores were treated as fixed effects, while the CHWs' organizations were treated as a random effect. We examined the interaction between time (pre- and post-program) and organization to assess consistency. A significant interaction would indicate variability in improvements across organizations, suggesting that data from all training programs should not be pooled for further analyses. Conversely, a non-significant interaction would support pooling the data for a more comprehensive analysis.

Quality Assurance Process

A process was developed to support CHWs in their new roles after completing in-person training. As CHWs enrolled eligible clients in the MTM service, they gathered patients' current medication lists from physicians and pharmacy prescription refill information. CHWs then met with patients and documented medication use, adherence barriers, and other relevant information. An experienced student pharmacist reviewed this information before sending to a partnering pharmacist. This ensured ongoing quality assurance and continuous improvement of CHWs' skills. Partnering pharmacists developed care plans based on the information shared by CHWs. CHWs used these care plan recommendations to resolve adherence barriers by interacting directly with patients and consulting with their medical providers as needed. About four weeks later, CHWs followed up with patients to check on the resolution of adherence barriers. This process was repeated as long as patients continued their relationship with CHWs.

Additionally, three-hour remediation training was provided for CHWs about three to four months after the live training program. All trained CHWs were invited to participate in the remediation session, where they met in small groups to practice the MTM process on a simulated patient case, discuss challenges,

and share strategies for resolving them. Documentation performed by CHWs on the simulated patient was assessed for accuracy and completeness.

Results

The training programs were offered in either English or Spanish, based on the CHWs' preferred language. A total of 109 CHWs participated in an in-person, live two-day training program format, which was preceded by six hours of pre-recorded video lectures.

A linear mixed-effects model was used to assess whether score improvements from pre- to post-program were consistent across the five organizations employing CHWs. The model revealed a significant main effect of time, with post-program scores being significantly higher than pre-program scores ($p < 0.001$), indicating an overall improvement in performance after training. The random effect of organization was small, and there was no significant interaction between time and organization, suggesting that the magnitude of improvement did not differ significantly between the five training programs. Therefore, CHWs from all five organizations experienced similar improvements, justifying the pooling of data across the different training programs for subsequent analyses.

The average pre-test score on the CHW MTM principles knowledge assessment for the 109 CHWs was 28.2%, which increased to 81.2% post-training. The average pre-test to post-test change score was 53% ($p < 0.001$).

Knowledge about Medication Therapy Management (MTM) Principles

CHWs showed substantial improvements across all thirteen MTM principles questions post-training, with significant differences in scores for questions related to the Drug Adherence Workup (DRAW)

tool,²⁹ Situation, Background, Assessment, Recommendation (SBAR) communication,³⁰ understanding of Proportion of Days Covered (PDC), interview strategies, medication history, utilizing open-ended questions, assessing pharmacy claims data (refill records), follow-up information, medical record history, medication adherence, and sharing expertise (see Table 2). These improvements remained statistically significant after applying the Bonferroni correction for multiple comparisons, indicating the robustness of the training program's impact.

Knowledge about Health Disparities and Cultural Competency and Self-Efficacy

The average pre-test score on the Knowledge about Health Disparities and Cultural Competency assessment was 76%, which increased to 78% post-training. Self-efficacy scores showed significant improvement across all 14 items measured, with all CHWs reporting high confidence in their ability to perform all 14 skills post-training (see Table 3). Pre-training self-efficacy scores ranged from 2.67 to 5.30 on a scale of 0 to 10, with the highest self-efficacy in asking the right questions to identify medication use practices for prescribed medications (Mean = 5.30) and the lowest in explaining the MTM program to others within the CHWs' organization (Mean = 2.67). The mean improvement from pre-training to post-training ranged from 3.59 to 5.57, with the greatest improvement observed in explaining the MTM program to others within the organization (Mean Improvement = 5.57). The percentage of CHWs who felt highly confident (scoring at least 7 out of 10) in performing each skill at the end of the training program ranged from 86% to 97%, indicating that most CHWs were highly confident in performing the 14 skills (see Table 3).

Quality Assurance

The quality assurance process where student pharmacists reviewed CHWs documentation following interactions with actual patients was helpful in identifying gaps in CHWs skills in performing

medication adherence patient workups. Every CHW had at least one deficiency cited by the student pharmacist. Examples of deficiencies included poor documentation, insufficient probing of the patient to uncover specific adherence barriers or the use of a self-medication such as herbal medications, or insufficient probing of the patient to uncover detailed information about how patients take medications daily. Uncovering these gaps led to one-on-one tutorials to address areas of concern. Student pharmacists monitored CHWs in subsequent patient encounters for improvement.

Remediation Training

Eighty-seven of the CHWs completed remediation training three to four months following the live training program, where they met in small groups to practice the MTM process on a simulated patient case. Sixty-eight of the CHWs correctly documented or performed in all skill areas and the remainder correctly documented or performed more than 80 percent of the tasks.

Discussion

This study demonstrates the effectiveness of a structured training program in improving CHWs' knowledge and self-efficacy in medication adherence support. The significant improvements in knowledge scores and self-efficacy suggest that the training successfully prepared CHWs with the necessary skills to address medication adherence issues in their communities and collaborate with the healthcare team. Our findings build upon previous research about CHW training programs related to medication management. For example, Jam et al.³¹ shared findings from a cross-sectional survey of 77 CHWs, which assessed CHW confidence and training gaps in medication management. They found that only 18 percent reported having received training about medication adherence and that more than half of CHWs said their confidence in providing medication management services was either poor or fair. Our

literature review found few reports of CHW training programs specifically focused on medication adherence, with most involving pharmacy technicians who had been cross-trained as CHWs.³²⁻³⁵

Our approach of combining pre-recorded lectures with live case-based interactive sessions taught within small groups of CHWs aligns with best practices in adult learning theory and mirrors successful approaches used in other health professional training programs. This blended learning approach has been increasingly recognized as effective in healthcare education.³⁶ Liu et al.³⁷ conducted a systematic review of blended learning in health professions, finding that it was more effective than or at least as effective as non-blended instruction for knowledge acquisition in health professions.

The use of case-based learning in small groups is supported by Thistlethwaite et al.,³⁸ who found in their systematic review that case-based learning is enjoyed by students and teachers and leads to improved learning outcomes in terms of clinical reasoning and knowledge. This approach is particularly relevant for CHWs, as it allows them to apply theoretical knowledge to practical, real-world scenarios they might encounter in their communities. Moreover, our focus on interactive, hands-on training aligns with the findings of Bluestone et al.,³⁹ who found that interactive methods, including case-based learning and hands-on practice sessions, were more effective than traditional didactic methods alone. The incorporation of pre-recorded lectures as part of a flipped classroom model is supported by research from Hew and Lo,⁴⁰ who found that flipped classrooms were either more effective than or as effective as traditional classrooms in terms of academic performance and student satisfaction.

Given their prior training and experience, it was expected that CHWs would perform well on pre-program assessments of health disparities and cultural competence. However, their post-program assessment scores did not show significant improvement. This could be attributed to the broader scope

of the questions, which focused on general health disparities and cultural competence rather than the specific cultural beliefs and values about medications covered in our training program. Additionally, the minimal improvement in scores may indicate that CHWs already possessed a relatively high level of competence in these areas before the training began.

The inclusion of cultural competency and health literacy in our training addresses a critical need identified in urban ambulatory settings, particularly for hypertension care among underserved populations. Mueller et al.⁴¹ demonstrated that culturally competent care can significantly improve blood pressure control in minority populations. Furthermore, Jolly et al.⁴² emphasized the role of health literacy in patient engagement and self-management of chronic conditions like hypertension. By equipping CHWs with these skills, we aimed to bridge the gap in care for underserved populations. This approach aligns with findings from Kangovi et al.,⁴³ who showed that CHWs trained in cultural competency and health literacy can improve chronic disease management. Integrating these competencies in CHW training also addresses the broader social determinants of health, as outlined by Marmot,⁴⁴ who argued that cultural factors and health literacy are crucial in reducing health inequities. Covert et al.⁴⁵ proposed a set of core competencies for Community Health Workers, and a focus on medication adherence in conjunction with health disparities would support the disease prevention and management competency. By focusing on these aspects, our training program aims to improve individual patient outcomes and contribute to the larger goal of health equity in diverse communities.

Our quality improvement process, involving ongoing support and feedback from student pharmacists and training remediation, ensures continuous learning and improvement beyond initial training, emphasizing the importance of structured support and iterative improvement in CHW training.⁴⁶ Moreover, our continuous improvement approach reflects best practices identified by Kok et al.,⁴⁷ emphasizing the need for ongoing supportive supervision and performance appraisal to maintain and

improve CHW effectiveness. Allen et al.⁴⁸ also demonstrated that continuous quality improvement initiatives lead to better health outcomes and increased job satisfaction among CHWs. Our experience shows that this process is essential for identifying training gaps and addressing deficiencies in critical skills for medication adherence patient workups.

Limitations and Future Directions

While we are confident in the training program's efficacy in improving knowledge and confidence among CHWs, some challenges were uncovered. Some CHWs did not fully engage with the at-home self-study materials before attending the in-person program. To address this, we implemented a short quiz on the subject matter before the training sessions, allowing us to identify and assist those who performed poorly. Further, some CHWs struggled to master motivational interviewing based on their performance, as assessed by the facilitator using a rubric, on a simulated patient during the remediation training exercise and instead relied on their own learned experiences. The limited MI training was insufficient for some CHWs. Future training programs should offer more intensive motivational interviewing training for those CHWs who need it.⁴⁹ We did not reassess our primary measures of knowledge and self-efficacy at any point other than before and immediately following the training program to determine whether improvement in knowledge and self-efficacy were sustained. However, we did check on skill performance during the remediation training, which was held about three to four months following the training, which showed that all trainees correctly documented or performed in most skill areas.

The CHWs in the training program were experienced professionals, many with prior training in social determinants of health and cultural competence, and some held CHW certification in Florida. Our MTM training was designed to complement their existing knowledge and skills, not replace comprehensive

CHW training curricula, which is commonly taught over many weeks. Future studies should explore the optimal duration and format of such training programs and strategies for ongoing education and support for CHWs.

Conclusion

Our structured training program, through a multi-faceted evidence-based approach effectively improves CHWs' knowledge and confidence in supporting medication adherence for hypertension patients.

Acknowledgements

David Angaran, Golnoosh Alipour-Haris, and Rupal Aroza made substantive contributions to the conduct of the project.

References

1. Neiman AB, Ruppert T, Ho M, et al. CDC Grand Rounds: Improving Medication Adherence for Chronic Disease Management — Innovations and Opportunities. *MMWR Morb Mortal Wkly Rep.* 2017;66. DOI: <http://dx.doi.org/10.15585/mmwr.mm6645a2>
2. Bosworth HB, Granger BB, Mendys P, Brindis R, et al. Medication adherence: a call for action. *Am Heart J.* 2011;162(3):412-424. DOI: 10.1016/j.ahj.2011.06.007.
3. Singh P. Medication Adherence: Adjunct to Patient Outcomes. *J Pract Cardiovasc Sci.* 2017;3(1):8-10. DOI: 10.4103/jpcs.jpcs_10_17
4. Brown MT, Bussell JK. Medication adherence: WHO cares? *Mayo Clin Proc.* 2011;86(4):304-314. DOI: 10.4065/mcp.2010.0575.
5. Sabaté E, ed. *Adherence to Long-term Therapies: Evidence for Action.* World Health Organization; 2003.
6. Mongkhon P, Ashcroft DM, Scholfield CN, Kongkaew C. Hospital admissions associated with medication non-adherence: a systematic review of prospective observational studies. *BMJ Qual Saf.* 2018;27(11):902-914. DOI: 10.1136/bmjqs-2017-007453.
7. Luga A, McGuire M. Adherence and health care costs. *Risk Manag Health Policy.* 2014;7:35-44. DOI.org/10.2147/RMHP.S19801
8. Kini V, Ho PM. Interventions to improve medication adherence: a review. *JAMA.* 2018;320(23):2461-2473. DOI:10.1001/jama.2018.19271
9. McDonnell PJ, Jacobs MR. Hospital admissions resulting from preventable adverse drug reactions. *Ann Pharmacother.* 2002;36:1331-1336.
10. Roger VL, Go AS, Lloyd-Jones DM, et al. Heart disease and stroke statistics—2012 update: a report from the American Heart Association. *Circulation.* 2012;125:e2-e220.

11. Glader EL, Sjolander M, Eriksson M, Lundberg M. Persistent use of secondary preventive drugs decline rapidly during the first 2 years after stroke. *Stroke*. 2010;41(2):397-401.
12. Burnier M, Egan BM. Adherence in hypertension: a review of prevalence, risk factors, impact, and management. *Circ Res*. 2019;124(7):1124-1140. DOI:org/10.1161/CIRCRESAHA.118.313220
13. Costa FV. Compliance with antihypertensive treatment. *Clin Exp Hypertens*. 1996;18(3-4):463-472.
14. Cramer JA, Benedict A, Muszbek N, Keskinaslan A, Khan ZM. The significance of compliance and persistence in the treatment of diabetes, hypertension and dyslipidemia: a review. *Int J Clin Pract*. 2008;62(1):76-87.
15. Martin LT, Parker RM. Insurance expansion and health literacy. *JAMA*. 2011;306(8):874-875. DOI:10.1001/jama.2011.1212
16. Baker DW, Parker RM, Williams MV, Clark WS. Health literacy and the risk of hospital admissions. *J Gen Intern Med*. 1998;13(12):791-798.
17. Ndumele CD, Shaykevich S, Williams D, Hicks LS. Disparities in adherence to hypertensive care in urban ambulatory settings. *J Health Care Poor Underserved*. 2010;21(1):132-143.
18. Johnson VR, Jacobson K, Gazmararian J, Blake S. Does social support help limited-literacy patients with medication adherence? *Patient Educ Couns*. 2010;79(1):14-24.
19. Brunner R, Dunbar-Jacob J, Leboff MS, et al. Predictors of adherence in the Women's Health Initiative Calcium and Vitamin D Trial. *Behav Med*. 2009;34(4):145-155.
20. Joyner-Grantham J, Mount DL, McCorkle OD, Simmons DR, Ferrario CM, Cline DM. Self-reported influences of hopelessness, health literacy, lifestyle action and patient inertia on blood pressure control in a hypertensive emergency department population. *Am J Med Sci*. 2009;338(5):368-372.
21. Molloy GJ, Perkins-Porras L, Strike PC, Steptoe A. Social networks and partner stress as predictors of adherence to medication, rehabilitation attendance, and quality of life following coronary syndrome. *Health Psychol*. 2008;27(1):52-58.

22. Segal R, Angaran DM, Odedina FT, Zeigler ML, Wallace JL. Opportunities and responsibilities for pharmacists to improve their effectiveness in addressing medication adherence through culturally sensitive collaborations with community health workers. *J Am Pharm Assoc* (2003). 2020;60(4):e25-e30. DOI.org/10.1016/j.japh.2020.02.023
23. Arvey SR, Fernandez ME. Identifying the core elements of effective community health worker programs: a research agenda. *Am J Public Health*. 2012;102(9):1633-1637. DOI: 10.2105/AJPH.2012.300649.
24. Brownstein JN, Bone LR, Dennison CR, Hill MN, Kim MT, Levine DM. Community health workers as interventionists in the prevention and control of heart disease and stroke. *Am J Prev Med*. 2005;29(5 Suppl 1):125-133.
25. Rosenthal EL, Brownstein JN, Rush CH, Hirsch GR, Willaert AM, Scott JR, Holderby LR, Fox DJ. Community health workers: part of the solution. *Health Aff (Millwood)*. 2010;29(7):1338-1342. DOI: 10.1377/hlthaff.2010.0081.
26. Brownstein JN, Chowdhury FM, Norris SL, Horsley T, Jack L, Zhang X, Satterfield D. Effectiveness of community health workers in the care of people with hypertension. *Am J Prev Med*. 2007;32(5):435-447.
27. Wheat L, Roane TE, Connelly A, et al. Using a Pharmacist-Community Health Worker Collaboration to Address Medication Adherence Barriers. *J Am Pharm Assoc* (2003). 2020;60(6):1009-1014.
28. Prieto-Merino D, Mulick A, Armstrong C, et al. Estimating proportion of days covered (PDC) using real-world online medicine suppliers' datasets. *J Pharm Policy Pract*. 2021;14:113.
29. Doucette WR, Farris KB, Youland KM, Newland BA, Egerton SJ, Barnes JM. Development of the Drug Adherence Work-up (DRAW) tool. *J Am Pharm Assoc* (2003). 2012;52(6):e199-e204. DOI: 10.1331/JAPhA.2012.12001

30. Agency for Healthcare Research and Quality. Tool: SBAR. Content last reviewed November 2019.
Available at: <https://www.ahrq.gov/sbar>.
31. Jam VA, McKay VA, Holmes JT. Identifying medication management confidence and gaps in training among community health workers in the United States. *J Community Health*. 2019;44(6):1180-1184. doi.org/10.1007/s10900-019-00688-9
32. Bailey JE, Surbhi S, Bell PC, Jones AM, Rashed S, Ugwueke MO. SafeMed: Using pharmacy technicians in a novel role as community health workers to improve transitions of care. *J Am Pharm Assoc (2003)*. 2016;56(1):73-81. DOI: 10.1016/j.japh.2015.11.011.
33. Yoon HS, Teshome BF, Eisenbeis A, Micek ST. Pharmacy technicians trained as community health workers: a prospective multicenter cohort study. *J Am Pharm Assoc (2003)*. 2024
34. Triyasakorn K, Nguyen E. Cross-training pharmacy personnel as community health workers. *J Am Pharm Assoc (2003)*. 2024;64(1):39-42.
35. Somerville P, Lindsay R, Nguyen E. Cross-Training Pharmacy Professionals as Community Health Workers. *Pharmacy (Basel)*. 2023;11(5):135.
36. Bosworth HB, Powers BJ, Olsen MK, et al. Home Blood Pressure Management and Improved Blood Pressure Control: Results From a Randomized Controlled Trial. *Arch Intern Med*. 2011;171(13):1173-1180. DOI:10.1001/archinternmed.2011.276
37. Liu Q, Peng W, Zhang F, et al. The Effectiveness of Blended Learning in Health Professions: Systematic Review and Meta-Analysis. *J Med Internet Res*. 2016;18(1):e2. DOI:10.2196/jmir.4807.
38. Thistlethwaite JE, Davies D, Ekeocha S, et al. The effectiveness of case-based learning in health professional education. A BEME systematic review: BEME Guide No. 23. *Med Teach*. 2012;34(6):e421-444. DOI: 10.3109/0142159X.2012.680939

39. Bluestone J, Johnson P, Fullerton J, et al. Effective in-service training design and delivery: evidence from an integrative literature review. *Hum Resour Health*. 2013;11:51. doi:10.1186/1478-4491-11-51.
40. Hew KF, Lo CK. Flipped classroom improves student learning in health professions education: a meta-analysis. *BMC Med Educ*. 2018;18:38. DOI.org/10.1186/s12909-018-1144-z
41. Mueller M, Purnell TS, Mensah GA, Cooper LA. Reducing racial and ethnic disparities in hypertension prevention and control: What will it take to translate research into practice and policy? *Am J Hypertens*. 2015;28(6):699-716.
42. Jolly SE, Navaneethan SD, Schold JD, et al. Development of a chronic kidney disease patient navigator program. *BMC Nephrol*. 2018;19(1):1-8.
43. Kangovi S, Mitra N, Grande D, et al. Patient-centered community health worker intervention to improve posthospital outcomes: A randomized clinical trial. *JAMA Intern Med*. 2017;177(7):905-913.
44. Marmot M. Health equity in England: the Marmot review 10 years on. *BMJ*. 2020;368:m693. doi:10.1136/bmj.m693. DOI:10.1136/bmj.m693
45. Covert H, Sherman M, Miner K, Lichtveld M. Core Competencies and a Workforce Framework for Community Health Workers: A Model for Advancing the Profession. *Am J Public Health*. 2019 Feb;109(2):320-327.
46. O'Brien MJ, Squires AP, Bixby RA, Larson SC. Role development of community health workers: An examination of selection and training processes in the intervention literature. *Am J Prev Med*. 2009;37(6 Suppl 1):S262-S269.
47. Kok MC, Dieleman M, Taegtmeier M, et al. Which intervention design factors influence performance of community health workers in low-and middle-income countries? A systematic review. *Health Policy Plan*. 2017;32(9):1207-1227.

48. Allen CG, Brownstein JN, Satsangi A, Escoffery C. Community Health Workers as Allies in Hypertension Self-Management and Medication Adherence in the United States, 2014. *Prev Chronic Dis.* 2016;13:160236. doi.org/10.5888/pcd13.160236
49. Berger BA, Villaume WA. *Motivational Interviewing for Health Care Professionals: A Sensible Approach.* 2nd ed. Washington, DC: APhA Press; 2020.

Journal Pre-proof

Figure 1 - Two Day CHW MTM Live Training Program Curriculum

Pre-In Class Homework Pre-Recorded Lecture Videos:

1. Pre-assessment
2. Watch video: CHW Introduction of the program including the Why in terms of Medication Safety
3. Watch video: Cultural Competency and Health Disparities
4. Watch video: Effective Communication and Motivational Interviewing

Day 1:

Time	Topic	Learning Objectives
7:30-8:00AM	Breakfast & Registration Sign-In	
8:00 – 8:15AM (15 Min)	Introductions & Expectations for program	Objectives: <ul style="list-style-type: none"> • Brief introduction of Facilitators (5 min) • Purpose & overview of training program (5 min) • Expectations and Requirements of Sponsoring Organization (5 min)
8:15-8:45AM (30 min)	MTM CHW process: Step by Step Practice Model	Objectives: <ul style="list-style-type: none"> • Why do we need a better solution • Describe each major step • CHW Role and Do/Don't • Pharmacist role • Practice Model of CHW and Pharmacist
8:45-9:30AM (45 Min)	The CMR (Comprehensive Medication Review): Personal Med List	Objectives: <ul style="list-style-type: none"> • What is a Personal Medication List (PML) and why is PML important to conducting an Adherence Workup and CMR • Establishing a Medication List • Discuss Current Medication List provided by Physician(s) and tactics for collecting information from a physician's office • Discuss Rx refill history and tactics for collecting refill history from a pharmacy • Rx history data sources; strengths, weaknesses and gaps • How to create a dialog with patient to learn about Rx and self-medications for establishing a medication list • Level of specificity needed about use of each Rx and Self-Medication within PML, including time of day, proximity to meals, dosage regimen on encounter form • Identifying discrepancies between Physician List, Rx refill history and Patient interview and how to document

9:30-10:00AM (30 Min)	Encounter Form Process	Objectives: <ul style="list-style-type: none"> • Introduce form • Review different sections • Introduce Jessica Faith patient case & how to use form
10:00-11:30AM (90 Min)	Medication List Breakout sessions	Objectives: <ul style="list-style-type: none"> • Review Jessica Faith PML • Creating a medication list from physician, pharmacy and patient sources • Practice asking patients questions and probing about meds including Self-medications • Documentation
11:30-12:15PM (45 Min)	Hypertension and Diabetes Review	Objectives: <ul style="list-style-type: none"> • Review topics about Understanding the disease, consequences of poorly controlled HTN and T2DM, Complications, and Treatment strategies • Understand medical management issues relevant to adherence workup in HTN and T2DM • Identifying Lifestyle issues HTN and T2DM • Emphasize what questions CHW should ask
12:15-12:45PM (30 Min)	Pharmacy ABC's	Objectives: <ul style="list-style-type: none"> • Walk in the shoes of a busy community pharmacist: competing pressures faced by a pharmacist on daily basis • Pharmacy types • Drug Tiers and Retail Cost sharing • Pharmacy Record Systems and how to understand Refill records
12:45-1:15PM	30 min Lunch Break	
1:15-1:45PM (30 Min)	Communication Motivational Interviewing (MI)	Objectives: <ul style="list-style-type: none"> • Understanding how MI is different from how most people communicate with one another • Applying MI to: (a) Word choice and vocabulary, (b) Patient recruitment, (c) Addressing non-adherence • HTN and Diabetes MI examples • Build rapport with patient and address issues they may have
1:45-2:00PM (15 Min)	MI Videos	Objectives: <ul style="list-style-type: none"> • Play videos (6.5 min) • 2nd video about (8 min)
2:00-2:15PM (15 Min)	MI Personal Story	Objectives: <ul style="list-style-type: none"> • Impact and strength MI can have
2:15-2:45PM (30 Min)	MI practice small groups	Objectives: <ul style="list-style-type: none"> • Practice MI case
2:45-3:30PM (45 Min)	Medication Adherence	Objectives: <ul style="list-style-type: none"> • Prevalence of adherence issues in HTN and T2DM • Most common adherence issues in HTN and T2DM • Alternate ways to uncover adherence issues: Rx

		refill history, Objective measures such as drug levels, patient self-report, patient interview <ul style="list-style-type: none"> • Assessing Health Literacy and why it's important • Discuss DRAW Adherence questions on form • Listening for Patient Concerns and Patient's self-reported Goals • Documenting adherence barriers
3:30-5:00PM (90 Min)	Medication Adherence Breakout sessions	Objectives: <ul style="list-style-type: none"> • Jessica Faith Case: Asking questions, active listening and probing • Practice adherence questions from encounter form without asking them directly • Identify the medication related problems (MRP) to be tackled • Documentation

Day 2:

Time	Topic	Presenter
7:30-8:00am	Breakfast & Sign-In	
8:00-8:30AM (30 Min)	Review Day-1 Topics	Objectives: <ul style="list-style-type: none"> • Review important training objectives from yesterday • CMR • Encounter Form • MI
8:30-9:00AM (30 Min)	Medication Adherence Review	Objectives: <ul style="list-style-type: none"> • Discuss set of Adherence questions on form • Listening for Patient Concerns and Patient's self-reported Goals • Documenting adherence barriers • Show a Poor Interview and an Excellent Interview
9:00-9:45AM (45 Min)	Care Plan and Documentation	Objectives: <ul style="list-style-type: none"> • Learning to be clear, concise when speaking with Patient • Communicating with physicians or pharmacists about MRPs

9:45-11:00AM (75 Min)	Care Plan Breakout Session -small groups	Objectives: <ul style="list-style-type: none"> • Jessica Faith Case: Practice discussing with a pharmacist uncovered MRPs and prioritizing them • Medication reconciliation and finalizing PML • Construct care plan/action plan • Practice discussion with Doctor • Documentation on Encounter form
11:00-11:45AM (45 Min)	Encounter form Process & Flow	Objectives: <ul style="list-style-type: none"> • Discuss logistics – sending forms to UF, communicating with MTM Center pharmacist, implementing MAP • Discuss follow-up encounters
11:45-12:30PM (45 Min)	Lunch break	
12:30-2:30PM (120 Min)	Lynda Blake Case Breakout session:	<ul style="list-style-type: none"> • Apply all lessons to the Lynda Blake Case
2:30-3:15PM (45 Min)	Debriefing – Review important Topics	
3:15-4:30PM (75 Min)	Post-assessment	

Table 1: Knowledge and Self-Efficacy Components of Pre- and Post-Test Training Material

Assessment	Components	Number of Questions	Type of Question	Number of points per Question	Total Points Possible
Knowledge Assessment of Principles MTM	<ul style="list-style-type: none"> • Principles of MTM support services • Scope of the CHW within MTM support services • Documentation Requirements • Gathering information pertinent to medications from 	13	Open-ended/ short answer	5	65 points possible and scaled on a range from 0 to 100%

	<p>various resources – the patient, prescription refill records, and the medical record</p> <ul style="list-style-type: none"> • Identification of medication adherence barriers • Communication skills, including motivational interviewing 				
Knowledge Assessment	<ul style="list-style-type: none"> • Health Disparities and Cultural Competence 	8	Multiple Choice	1	8 points possible and scaled on a range from 0 to 100%
Self-Efficacy of Providing MTM Services	<ul style="list-style-type: none"> • Ability to identify and enroll patients • Provide MTM support services • Document services 	14	Scale ranging from 0 (cannot do at all) to 10 (highly certain can do)	<p>Each item scored from 0 to 10 by respondent. Each response assigned to a category according to the following ranges:</p> <p>Low self-efficacy (0 to 3);</p> <p>Moderate self – efficacy (4 to 6);</p> <p>High self-efficacy (7 or greater)</p>	Reported as improvement from pre- to post-training and as percentage of trainees falling within each range of self-efficacy (low, moderate, high)

Table 2: Performance on MTM Practices Knowledge Assessment

MTM Practices Knowledge Question	Pre-Training Mean Score ^a	Post-Training Mean Score ^a
1. What are the questions in the DRAW tool and the suggested response?	0.0±0.0	5.0±0.0 ^b
2. When examining a medication profile and refill history, what are the red flags to look for?	2.3±1.0	4.0±0.9 ^b
3. What does SBAR ^c stand for regarding communicating with other healthcare professionals?	1.7±0.8	4.9±0.2 ^b
4. What does the PDC ^c tell you and not tell you?	0.4±0.3	4.6±0.5 ^b
5. What is an interview strategy to help patients remember the OTCs, herbal, dietary supplements and home remedies they use?	0.3±0.2	3.8±0.8 ^b
6. What are the strengths and weaknesses of a patient-provided medication history?	2.4±0.8	4.6±0.9 ^b
7. Provide an example of an open-ended “why” question without using the word why.	1.6±0.7	4.3±0.8 ^b
8. What contribution to your developing a personal medication list and care plan would you expect a participating pharmacist to do?	1.1±0.9	2.4±0.8 ^b
9. What are the strengths and weaknesses of a pharmacy refill record?	2.3±0.8	4.7±0.6 ^b
10. What information would you gather during a follow-up to the initial CMR ^c interview?	1.7±0.9	3.3±0.8 ^b
11. What are the strengths and weaknesses of a patient’s medical record base/ medication history?	2.7±0.8	4.7±0.4 ^b
12. List the ways to determine medication adherence.	1.6±1.0	3.4±0.9 ^b
13. List the four key steps to sharing expertise with the patient.	0.0±0.0	3.1±0.8 ^b

^a Scores range from 0 (poor response) to 5 (excellent response)

^b Post-training score greater than pre-training score (p<0.0038)

^c Drug Adherence Workup (DRAW) tool, Situation, Background, Assessment, Recommendation (SBAR) communication, Proportion of Days Covered (PDC), Comprehensive Medication Review (CMR).

Table 3. Self-Efficacy for Activities/ Tasks Related to MTM

Self-Efficacy Item	Mean Pre-training \pm Standard Deviation	Mean Change Score \pm Standard Deviation	Percent Post Score Highly Confident ^b
ID & Enroll Patients			
Identify a Patient that is eligible for MTM Service	3.65 \pm 3.17	5.19 \pm 3.74 ^a	95
Explain the purpose of the MTM Service to a patient	3.73 \pm 3.73	5.35 \pm 3.75 ^a	95
Motivate a Patient who is not interested in MTM Service	3.40 \pm 3.20	4.84 \pm 3.53 ^a	92
Provide Services			
Gather necessary patient-specific data from the treating physicians for MTM encounter form	4.05 \pm 3.16	4.62 \pm 3.69 ^a	92
Gather necessary prescription data from the pharmacy for MTM encounter form	4.43 \pm 3.43	5.24 \pm 3.65 ^a	95
Determine the level of literacy using a validated system	4.24 \pm 3.41	3.59 \pm 3.58 ^a	89
Ask the right questions to identify medication use practices for prescribed medications	5.30 \pm 3.52	4.84 \pm 3.22 ^a	97
Ask the right questions to uncover barriers that affect the use of prescribed medications	4.35 \pm 3.12	4.30 \pm 3.42 ^a	95
Ask the right questions to identify medication use practices involving self-medication	4.89 \pm 3.43	4.32 \pm 3.51 ^a	95
Perform a medication reconciliation to establish a complete and accurate medication list	4.89 \pm 3.39	4.27 \pm 3.96 ^a	92
Apply motivational interviewing techniques to improve medication use practices	4.72 \pm 3.37	4.46 \pm 3.23 ^a	92
In collaboration with the pharmacist, verbally communicate recommendations to a patient	4.40 \pm 3.27	3.76 \pm 3.41 ^a	97
Explain MTMS to others within your organization	2.67 \pm 2.94	5.57 \pm 3.19 ^a	86
Document Services			
Appropriately document information on the encounter form for pharmacists to review	4.54 \pm 3.35	5.24 \pm 3.12 ^a	92

^aImprovement in self-efficacy from pre-training to post-training periods was statistically significant ($p < 0.0002$), based on analysis of pair t-tests

^bHighly confident was a score of at least 7 on a scale ranging from 0 (cannot do at all) to 10 (highly certain can do)

CRedit author statement: Evaluating a Community Health Worker Training Program for Medication Adherence Support

Richard Segal: Conceptualization, Methodology, Formal Analysis, Investigation, Writing – Original Draft, Writing – Review & Editing, Supervision, Project administration, Funding Acquisition. **Michelle L. Zeigler:** Investigation, Writing – Original Draft, Writing – Review & Editing, **Jordan L. Wallace:** Investigation, Writing – Original Draft, Writing – Review & Editing, Project Administration, **Teresa E. Roane:** Investigation, Writing – Review & Editing, **Lorraine Mobley:** Investigation, Writing – Review & Editing, Folakemi T. Odedina: Conceptualization, Investigation, Writing – Review & Editing.

Journal Pre-proof